

**Table. Distribution of admissions for acute metabolic decompensation (coma) and micro- and macrovascular complications among patients with type 1 and type 2 diabetes, Emilia-Romagna, 2000**

ICD9-CM (Primary diagnosis)	Rate Type		Tasso Tipo		Ratio of rates		
	Diabetes Type 1 N=3,044	1 (x 1000 exempt)	Diabetes Type 2 N= 17,448)	2 (x 1000 exempt)	Type 2/Type 1	95% CI)	
Ketoacidotic coma	269	88.4	382	21.9	0.25	(0.21	÷ 0.29)
Kidney disease	129	42.4	376	21.5	0.51	(0.42	÷ 0.62)
Circulatory disturbances	271	89.0	1039	59.5	0.67	(0.59	÷ 0.76)
Gangrene	95	31.2	173	9.9	0.32	(0.25	÷ 0.41)
Peripheral vasculopathy	41	13.5	108	6.2	0.46	(0.32	÷ 0.66)
Heart failure	268	88.0	1823	104.5	1.19	(1.05	÷ 1.34)
Myocardial infarct	187	61.4	1362	78.1	1.27	(1.10	÷ 1.47)
TIA* o stroke	110	36.1	887	50.8	1.41	(1.16	÷ 1.71)
Eye findings	76	25.0	483	27.7	1.11	(0.87	÷ 1.41)
Neurologic problems	66	21.7	308	17.7	0.81	(0.63	÷ 1.06)
Hyperosmolar coma	26	8.5	139	8.0	0.93	(0.61	÷ 1.42)

95% CI= 95% confidence interval

\* TIA = transient ischemic attack