



Ospedali Riuniti di Bergamo



Roma, 10 dicembre 2012

A Giusti, ML Farina, A Clavenna & Gruppo di lavoro Farfalla

Benefici dell'allattamento al seno



Breastfeeding is the ideal way of providing young infants with the nutrients they need for healthy growth and development.

Virtually all mothers can breastfeed, provided they have accurate information, and the support of their family and the health care system.

Exclusive breastfeeding is recommended up to 6 months of age.

Thereafter infants should receive complementary foods with continued breastfeeding up to 2 years of age or beyond.



Keywords



... provided they have

accurate information,
and the support of their family and
the health care system



United States
National Library
of Medicine

• LactMed

2

Paroxetine
CASRN: 61869-08-7

Drug Levels and Effects:

Summary of Use during Lactation:

Because of the low levels of **paroxetine** in breastmilk, amounts ingested by the infant are small and **paroxetine** has not been detected in the serum of any infant tested. ...

... Most authoritative reviewers consider **paroxetine** one of the preferred antidepressants during breastfeeding.

Accurate
information



Farmaci e allattamento

Accurate information



L' 80% dei foglietti informativi riporta
la controindicazione all'uso
in gravidanza e allattamento,
anche quando
la sicurezza di impiego del farmaco
è documentata



Farmaci e allattamento

Scarsità di informazioni attendibili



scelta tra curarsi o allattare



sospensione o interruzione precoce
dell'allattamento

Quasi tutti i farmaci
vengono escreti nel latte materno,
ma questo non comporta necessariamente
un rischio per il lattante

TIS Bergamo


gravidanza

- ✓ Stima del rischio di teratogenesi
- ✓ opportunità diagnostica prenatale
- ✓ identificazione di farmaco/i con stessa indicazione terapeutica e minor rischio
- ✓ follow-up della gravidanza


allattamento

- ✓ compatibilità tra farmaco e allattamento
- ✓ identificazione farmaco/i con stessa indicazione terapeutica e maggior sicurezza per il lattante

**Centro Antiveleni (CAV)
Servizio d'Informazione Teratologica (TIS)
anno 2011
totale consulenze = 20.400**



Centro Antiveneni (CAV)
Servizio d'Informazione Teratologica (TIS)
anno 2011
totale consulenze TIS = 12.358



... le donne sono sole?



2011 – allattamento 6.422 consulenze




stima del rischio

Allattamento → 6.422 consulenze



compatibile
71%



Uso
e

C

F



The logo is contained within a light blue rounded rectangle with a green border. At the top left, there is a stylized butterfly with green and purple wings. Below it is a line drawing of a baby's head in profile, looking upwards. The word "FARFALLA" is written in large, purple, rounded letters across the middle. At the bottom, the text "Uso dei FARMaci, Farmacovigilanza e gestione dell'ALLAttamento" is written in a smaller, dark blue font.

anza
o

o

a

Attività



2012

- Collaborazione tecnico-scientifica multidisciplinare
- Pacchetto formativo costruito in forma partecipativa e validato
- Studio preliminare di conoscenze, atteggiamenti e pratiche

2013

- Sperimentazione di interventi comunicativi e formativi su diversi target

Obiettivi

- ✓ promozione della comunicazione e del lavoro in rete tra professionisti e tra servizi
- ✓ promozione dell'health literacy ed empowerment delle madri allattanti che devono ricorrere all'uso di farmaci

Risultati attesi

Maggiore conoscenza della
gestione dell'allattamento

Modifiche dei
comportamenti prescrittivi
in allattamento

grazie!

