

Table – Distribution of the 10 most common *Salmonella* serotypes isolated in Italy by source, ENTER-NET 2000

Serotype	Man		Animals		Food/ envirnoment		Unknown		Total	
	N.	%	N.	%	N.	%	N.	%	N.	%
<i>Typhimurium</i>	1,848	31.0	274	24.8	716	19.2	8	11.0	2,846	26.2
<i>Enteritidis</i>	2,459	41.2	50	4.5	241	6.5	7	9.6	2,757	25.4
<i>Infantis</i>	434	7.3	19	1.7	436	11.7	15	20.5	904	8.3
<i>Derby</i>	113	1.9	45	4.1	331	8.9	4	5.5	493	4.5
<i>Livingstone</i>	38	0.6	212	19.2	175	4.7	1	1.4	426	3.9
<i>Bredeney</i>	72	1.2	39	3.5	129	3.5	4	5.5	244	2.2
<i>Blockley</i>	61	1.0	42	3.8	99	2.7	4	5.5	206	1.9
<i>Anatum</i>	23	0.4	22	2.0	125	3.4	3	4.1	173	1.6
<i>Brandenburg</i>	102	1.7	5	0.5	54	1.5	1	1.4	162	1.5
<i>London</i>	31	0.5	10	0.9	111	3.0	0	0.0	152	1.4
Other	785	13.2	387	35.0	1,303	35.0	26	35.6	2,501	23.0
Total	5,966	100.0	1,105	100.0	3,720	100.0	73	100.0	10,864	100.0