



Characteristics of SARS-CoV-2 patients dying in Italy Report based on available data on January 27th, 2021

1. Sample

The present report describes the characteristics of 85,418 SARS-CoV-2 patients dying in Italy*, as reported by the Integrated Covid-19 Surveillance System coordinated by the National Institute of Health-ISS. Geographic distribution across the 19 regions and 2 autonomous provinces of Trento and Bozen is presented in *Table 1*. Absolute number and percentage of deaths are reported according to the 3 phases that characterized the pandemic from the beginning in 2020 to the 27th of January 2021: the first wave (March-May 2020), the low incidence phase (June-September 2020), and the second wave (October 2020 – January 2021), the latter is still ongoing.

The surveillance data on deaths are constantly updated and consolidated and both activities require time to be carried out. Each Region has its own organization for updating data and related execution times. Therefore, when reading the data by Region, it is necessary to take into account that the timing of notification, control, verification and updating of data varies from Region to Region and from period to period; this may involve variations (increasing and/or decreasing) and differences both with the data already published in the previous reports, and with the data published by the Civil Protection. It should also be considered that the regional differences in the percentage of deaths reported in the table should not be interpreted in terms of risk. In fact, case fatality depends on the number of infections occurring in each region over a period of time compatible with the possible observation of the fatal event. In this regard, it should also be taken into account that the difference between the number of infections and the notified cases may vary regionally and over time as a function of different test access strategies adopted during the epidemic, thus distorting the comparison of regional case fatality.

Table 1. Geographic distribution of deceased patients SARS-CoV-2 positive by period

REGION	March-May 2020		June-Sept. 2020		Oct. 2020- 27 th Jan. 2021		Total	
	N	%	N	%	N	%	N	%
Lombardia	16,362	47.7	608	33.1	9,704	19.7	26,674	31.2
Emilia-Romagna	4,313	12.6	173	9.4	4,782	9.7	9,268	10.9
Veneto	1,949	5.7	247	13.4	6,528	13.2	8,724	10.2
Piemonte	3,992	11.6	123	6.7	3,923	8.0	8,038	9.4
Lazio	840	2.5	144	7.8	3,747	7.6	4,731	5.5
Toscana	1,046	3.1	96	5.2	2,870	5.8	4,012	4.7
Campania	477	1.4	59	3.2	3,234	6.6	3,770	4.4
Sicilia	300	0.9	58	3.2	2,932	6.0	3,290	3.9
Liguria	1,516	4.4	138	7.5	1,558	3.2	3,212	3.8
Puglia	524	1.5	76	4.1	2,492	5.1	3,092	3.6
Friuli-Venezia Giulia	347	1.0	21	1.1	1,695	3.4	2,063	2.4
Marche	978	2.9	7	0.4	991	2.0	1,976	2.3
Abruzzo	446	1.3	37	2.0	956	1.9	1,439	1.7
Trento	402	1.2	4	0.2	733	1.5	1,139	1.3
Sardegna	131	0.4	24	1.3	790	1.6	945	1.1

Bolzano	290	0.8	2	0.1	586	1.2	878	1.0
Umbria	75	0.2	9	0.5	653	1.3	737	0.9
Calabria	96	0.3	4	0.2	375	0.8	475	0.6
Valle d'Aosta	143	0.4	3	0.2	257	0.5	403	0.5
Molise	22	0.1	2	0.1	238	0.5	262	0.3
Basilicata	29	0.1	2	0.1	230	0.5	261	0.3
Total	34,278	100.0	1,837	100.0	49,274	100.0	85,389	100.0

Note: For 29 deceased persons, period was not possible to be evaluated; the number of deaths, relating to the period October 2020-January 2021, is in the consolidation phase due to the delay in notification.

* SARS-CoV-2 related deaths presented in this report are those occurring in patients who test positive for SARS-CoV-2RT by PCR, independently from pre-existing diseases.

2. Demographics

Mean age of patients dying for SARS-CoV-2 infection was 81 years (median 83, range 0-109, IQR 75-88). Women were 37,295 (43.7%). *Figure 1* shows that median age of patients dying for SARS-CoV-2 infection was more than 30 years higher as compared with the national sample diagnosed with SARS-CoV-2 infection (median age 48 years). *Figure 2* shows the absolute number of deaths by age group. Women dying for SARS-CoV-2 infection had an older age than men (median age women, 86 years - median age men, 80 years).

Figure 1. Median age of patients with SARS-CoV-2 infection and SARS-CoV-2 positive deceased patients




Figure 2. Absolute number of deaths by age group





Figure 3 shows the trend in the average age of SARS-CoV-2 positive deceased patients per calendar week, starting from the 3rd week of February 2020 (the date of the first death dates back to 21st February 2020). The average age of weekly deceased persons has substantially increased up to 85 years (1st week of July) and then dropped slightly.

Figure 3. Mean age of SARS-CoV-2 positive deceased patients by week of death



3. Deaths under the age of 50 years

As of January 27th 2021, 941 out of the 85,418 (1,1%), positive SARS-CoV-2 patients under the age of 50 died. In particular, 234 of these were less than 40 years (138 men and 96 women), age range between 0 and 39 years. For 52 patients under the age of 40 years no clinical information is available; out of the remaining ones, 147 had serious pre-existing pathologies (cardiovascular, renal, psychiatric pathologies, diabetes, obesity) and 35 had no major pathologies.

4. Pre-existing conditions

Table 2 presents most common comorbidities diagnosed before SARS-CoV-2 infection in a sample of SARS-CoV-2 positive deceased patients. Data on diseases were based on chart review and was available on 6,381 patients dying in-hospital for whom it was possible to analyse clinic charts. The medical records are sent to the ISS by the hospitals at different times, compatibly with the priorities of the activities carried out in the hospitals themselves. Therefore, the sample is opportunistic; it represents deaths in subjects who needed hospitalization only, and the regions are represented trying to maintain a proportionality with respect to the number of deaths. Mean number of diseases was 3.6 (median 3, SD 2.1). Overall, 3.1% of the sample presented with a no comorbidities, 12.1% with a single comorbidity, 18.6% with 2, and 66.3% with 3 or more.

Before hospitalization, 21% of SARS-CoV-2 positive deceased patients followed ACE-inhibitor therapy and 13% angiotensin receptor blockers-ARBs therapy. This information can be underestimated because data on drug treatment before admission were not always described in the chart.

Table 2. Most common comorbidities observed in SARS-CoV-2 positive deceased patients

Diseases	N	%
<i>Ischemic heart disease</i>	1,790	28.1
<i>Atrial Fibrillation</i>	1,547	24.2
<i>Heart failure</i>	1,028	16.1
<i>Stroke</i>	733	11.5
<i>Hypertension</i>	4,200	65.8
<i>Type 2-Diabetes</i>	1,868	29.3
<i>Dementia</i>	1,501	23.5
<i>COPD (Chronic Obstructive Pulmonary Disease)</i>	1,112	17.4
<i>Active cancer in the past 5 years</i>	1,065	16.7
<i>Chronic liver disease</i>	309	4.8
<i>Chronic renal failure</i>	1,339	21.0
<i>Dialysis</i>	137	2.1
<i>Respiratory failure</i>	425	6.7
<i>HIV Infection</i>	16	0.3
<i>Autoimmune diseases</i>	267	4.2
<i>Obesity</i>	688	10.8
Number of comorbidities		
<i>0 comorbidities</i>	196	3.1
<i>1 comorbidity</i>	772	12.1
<i>2 comorbidities</i>	1,185	18.6
<i>3 comorbidities and over</i>	4,228	66.3

Table 3 presents the most common pre-existing chronic pathologies in patients who died, separately in men (n = 3,835) and women (n = 2,546). The average number of pathologies observed in women is 3.8 (median 4, Standard Deviation 2.0). In men the average number of pathologies observed is 3.5 (median 3, Standard Deviation 2.1).

Table 3. Most common comorbidities observed in SARS-CoV-2 positive deceased patients by gender

Diseases	Women		Men	
	N	%	N	%
<i>Ischemic heart disease</i>	587	23.1	1,203	31.4
<i>Atrial Fibrillation</i>	648	25.5	899	23.4
<i>Heart Failure</i>	464	17.8	564	14.5
<i>Stroke</i>	315	12.4	418	10.9
<i>Hypertension</i>	1,733	68.1	2,467	64.3
<i>Type 2-Diabetes</i>	684	26.9	1,184	30.9
<i>Dementia</i>	824	32.4	677	17.7
<i>COPD (Chronic Obstructive Pulmonary Disease)</i>	355	13.9	757	19.7
<i>Active cancer in the past 5 years</i>	386	15.2	679	17.7
<i>Chronic liver disease</i>	110	4.3	199	5.2
<i>Chronic renal failure</i>	492	19.3	847	22.1
<i>Dialysis</i>	42	1.6	95	2.5
<i>Respiratory failure</i>	176	6.9	249	6.5
<i>HIV Infection</i>	2	0.1	14	0.4
<i>Autoimmune diseases</i>	149	5.9	118	3.1
<i>Obesity</i>	275	10.8	413	10.8
Number of comorbidities				
<i>0 comorbidities</i>	50	2.0	146	3.8
<i>1 comorbidity</i>	272	10.7	500	13.0
<i>2 comorbidities</i>	447	17.6	738	19.2
<i>3 comorbidities and over</i>	1,777	69.8	2,451	63.9

Table 4 presents the most common pre-existing chronic diseases in deceased patients divided into 4 age groups (16-59, 60-69, 70-79, 80+ years). The prevalence of *ischemic heart disease*, *atrial fibrillation*, *heart failure*, *stroke*, *arterial hypertension*, *dementia*, *chronic renal failure*, *respiratory failure* increase with age. On the other hand, prevalence of *chronic liver disease*, diseases for which *dialysis* is required, *HIV infection*, and *obesity* decrease with age increasing; for *diabetes*, *COPD*, and *cancer* prevalence decreases only in the last age group in contrast to the growth with age; for *autoimmune diseases*, on the contrary, prevalence increases only in the last age group, in contrast to the decreasing with age. As for the number of pathologies, the prevalence of those with 3 or more pathologies increases with age, while the prevalence of those with less than 3 pathologies decreases with age. For all the considered pathologies, the *trend* is statistically significant

Table 4. Most common comorbidities observed in SARS-CoV-2 positive deceased patients by age-groups

Age-groups	16-59		60-69		70-79		80+		Total	
	(n=445)		(n=652)		(n=1,597)		(n=3,683)		(n=6,377)	
Diseases	N	%	N	%	N	%	N	%	N	%
Ischemic heart disease	32	7.2	138	21.2	468	29.3	1,152	31.3	1,790	28.1
Atrial Fibrillation	12	2.7	69	10.6	305	19.1	1,161	31.5	1,547	24.3
Heart Failure	22	4.9	67	10.3	216	13.5	722	19.6	1,027	16.1
Stroke	14	3.1	60	9.2	167	10.5	492	13.4	733	11.5
Hypertension	164	36.9	390	59.8	1,083	67.8	2,563	69.6	4,200	65.9
Type 2-Diabetes	104	23.4	220	33.7	567	35.5	977	26.5	1,868	29.3
Dementia	13	2.9	40	6.1	181	11.3	1,267	34.4	1,501	23.5
COPD (Chronic Obstructive Pulmonary Disease)	34	7.6	83	12.7	313	19.6	682	18.5	1,112	17.4
Active cancer in the past 5 years	74	16.6	125	19.2	330	20.7	535	14.5	1,064	16.7
Chronic liver disease	33	7.4	40	6.1	97	6.1	139	3.8	309	4.8
Chronic renal failure	49	11.0	96	14.7	310	19.4	884	24.0	1,339	21.0
Dialysis	19	4.3	18	2.8	45	2.8	55	1.5	137	2.1
Respiratory failure	20	4.5	27	4.1	96	6.0	282	7.7	425	6.7
HIV Infection	10	2.2	2	0.3	3	0.2	1	0.0	16	0.3
Autoimmune diseases	30	6.7	35	5.4	55	3.4	147	4.0	267	4.2
Obesity	130	29.2	142	21.8	206	12.9	210	5.7	688	10.8
Number of comorbidities										
0 comorbidities	46	10.3	41	6.3	50	3.1	59	1.6	196	3.1
1 comorbidity	115	25.8	121	18.6	217	13.6	318	8.6	771	12.1
2 comorbidities	103	23.1	150	23.0	321	20.1	610	16.6	1,184	18.6
3 comorbidities and over	181	40.7	340	52.1	1,009	63.2	2,696	73.2	4,226	66.3


5. Diagnosis of hospitalization

In 90.7% of hospitalized persons who died and whose medical records were analysed (N=6,381), conditions (e.g. pneumonia, respiratory failure) or symptoms (e.g. fever, dyspnoea, cough) compatible with SARS-CoV-2 were mentioned. In 554 cases (9.3% of cases) the diagnosis of hospitalization was not related to the infection. In 77 cases the diagnosis of hospitalization concerned exclusively neoplastic pathologies, in 184 cases cardiovascular pathologies (for example Acute Myocardial Infarction-AMI, heart failure, stroke), in 74 cases gastrointestinal pathologies (for example cholecystitis, perforation of the intestine, intestinal obstruction, cirrhosis), in 219 cases other pathologies.

6. Symptoms

Figure 4 shows symptoms most commonly observed at hospital admission of hospitalized persons who died and whose medical records were analysed (N=6,381). Fever, dyspnoea and cough were the most commonly observed symptoms, while diarrhoea and haemoptysis were less commonly observed. Overall, 8.1% of patients did not present any symptoms at hospital admission.

Figure 4. Most common symptoms observed in SARS-CoV-2 positive deceased patients



7. Acute conditions

Between patients whose medical records were analysed, Acute Respiratory Distress syndrome was observed in the majority of patients (94.0% of cases), followed by acute renal failure (24.4%). Superinfection was observed in 19.6% and acute cardiac injury in 10.8% of cases.

8. Treatments


Antibiotics were used by 85.9% of patients during hospital stay, while less used were corticosteroids (53.3%) and antivirals (46.4%). Concomitant use of these 3 treatments was observed in 23.1% of cases.

Out of SARS-CoV-2 positive deceased patients, 4.0% were treated with Tocilizumab during hospitalization.

9. Time-line

Figure 5 shows, for SARS-CoV-2 positive deceased patients whose medical records were analysed (N=6,381), the median times, in days, from the onset of symptoms to death (12 days), from the onset of symptoms to hospitalization (5 days) and from hospitalization to death (7 days). The time from hospitalization to death was 6 days longer in those who were transferred to intensive care than those who were not transferred (12 days vs. 6 days).

Figure 5. Median hospitalization times (in days) in SARS-CoV-2 positive deceased patients



10. Comparison of death characteristics in the 3 quarters March-May 2020, June- September 2020, and October 2020 – January 2021

Table 5 summarizes the main characteristics of deaths with COVID-19 that occurred in 3 periods of time from the beginning of the pandemic in 2020: the initial quarter, March-May 2020, the second quarter, June-September 2020, and the third period October 2020 – January 2021. Overall, the sample represents 7.6% of all deaths from the beginning of the pandemic; in particular, the 13.5% of those who died between March and May 2020, 28.5% of those who died between June and September 2020, and 2.8% of those who died between October 2020 and January 2021.

Deaths of people with 3 or more pre-existing pathologies increase and those of persons with fewer pathologies or none decrease: this seems to indicate that in the second and third periods deaths concern older people and persons with a pre-existing health condition worse than those dying in the first quarter (table 5).

The use of drugs is also extremely different in the three periods, with a clear reduction in the use of antivirals and an increase in the use of steroids in the second and third periods.

Table 5. Mean age, prevalence of women, number of pre-existing diseases, complications and treatments in deaths with COVID-19 in the 3 periods March-May 2020, June-September 2020, and October 2020–January 2021

Sample of the evaluated clinical charts	All (n=6,381)	March-May 2020 (n=4,532)	June-Sept. 2020 (n=518)	Oct. 2020 – January 2021 (n=1,331)	<i>p-value*</i>
Mean age (years)	79.1	78.5	81.1	80.7	<0.001
	n (%)	n (%)	n (%)	n (%)	
Women	2,546 (39.9)	1,743 (38.5)	267 (51.5)	536 (40.3)	<0.001
<i>N of comorbidities</i>					
0	196 (3.1)	169 (3.7)	8 (1.5)	19 (1.4)	<0.001
1	772 (12.1)	608 (13.4)	44 (8.5)	120 (9.0)	
2	1,185 (18.6)	915 (20.2)	70 (13.5)	200 (15.0)	
3 or more	4,228 (66.3)	2,840 (62.7)	369 (76.4)	992 (74.5)	
<i>Complications during hospitalization</i>					
Acute Respiratory Distress Syndrome	5,852 (94.0)	4,159 (95.1)	425 (83.3)	1,268 (94.2)	<0.001
Acute renal failure	1,517 (24.4)	1,003 (22.9)	140 (27.5)	374 (27.8)	<0.001
Acute cardiac injury	671 (10.8)	472 (10.8)	51 (10.0)	148 (11.0)	0.823
Superinfection	1,218 (19.6)	728 (16.7)	215 (42.2)	275 (20.4)	<0.001
<i>Treatments</i>					
Antibiotics	5,378 (82.5)	3,830 (86.7)	438 (86.2)	1,110 (82.9)	0.002
Antivirals	2,903 (46.4)	2,606 (59.0)	162 (31.9)	135 (10.1)	<0.001
Steroids	3,341 (53.3)	1,911 (43.3)	333 (65.6)	1,097 (81.9)	<0.001
Tocilizumab	229 (4.0)	172 (4.4)	29 (5.9)	28 (2.2)	<0.001

* *p-value* for difference between the 3 periods

The distribution of the main pre-existing diseases in the different periods is presented in *figure 6*. The prevalence of atrial fibrillation, stroke, dementia, COPD, cancer, renal insufficiency, and obesity varies significantly in the three periods. These pathologies are more frequently diagnosed in the deceased in the second and third period than in the first (*Figure 6*).

Figure 6. Pre-existing pathologies in deaths with COVID-19 in the 3 periods




Table 6 shows the durations, as median times (in days), from the symptoms onset to death, SARS-CoV-2 testing, and hospitalization, and from the hospitalization to death, in the 3 periods considered. Between the first and second period doubles the time that passes from the onset of symptoms to death, while it returns to the initial levels in the third period; the time from the onset of symptoms to the swab for the detection of SARS-CoV-2 infection decreases in the second period and remain stable in the third, as well as the time between the onset of symptoms and hospitalization; the median duration in days from hospitalization to death doubles between the first and second period; it decreases again in the third period. These results seem to suggest a greater reactivity of the Health System evidenced by the greater speed in carrying out diagnostic tests and hospitalization.

Table 6. Median times (in days) between symptoms onset, PCR test, hospitalization and death in the 3 periods

Times(in days)	All (n=6,381)		March-May 2020 (n=4,532)		June-Sept. 2020 (n=518)		Oct. 2020 – January 2021 (n=1,331)		p- value*
	Median	IQR	Median	IQR	Median	IQR	Median	IQR	
From symptoms onset to death	12	7-20	12	7-19	23	10-57	12	7-19	<0.001
From symptoms onset to SARS-CoV-2 testing	4	2-8	5	2-9	3	0-7	3	0-6	<0.001
From symptoms onset to hospitalization	4	2-7	4	2-7	3	1-7	3	1-7	<0.001
From hospitalization to death	7	3-14	7	3-13	15	6-45	7	3-13	<0.001

* p-value for difference between the 3 periods

IQR = Inter-Quartile Range

The data here presented can be explained by a greater knowledge about the infection and a greater ability and timeliness of treatment in the period June-August in comparison to the previous quarter. In addition, it is likely that in the months of March and April SARS-CoV-2 infection was under-diagnosed in many frail elderly who died (such as those living in RSA). This may have led to an underestimation of the burden of the diseases in persons dying in that period

This report was produced by SARS-CoV-2 Surveillance Group

Members of the SARS-CoV-2 Surveillance Group

Luigi Palmieri, Elvira Agazio, Xanthi Andrianou, Pierfrancesco Barbariol, Antonino Bella, Eva Benelli, Luigi Bertinato, Matilde Bocci, Stefano Boros, Giovanni Calcagnini, Marco Canevelli, Federica Censi, Alessandra Ciervo, Elisa Colaizzo, Martina Del Manso, Corrado Di Benedetto, Chiara Donfrancesco, Massimo Fabiani, Francesco Facchiano, Marco Florida, Fabio Galati, Marina Giuliano, Tiziana Grisetti, Cecilia Guastadisegni, Yllka Kodra, Ilaria Lega, Cinzia Lo Noce, Pietro Maiozzi, Valerio Manno, Margherita Martini, Alberto Mateo Urdiales, Eugenio Mattei, Claudia Meduri, Paola Meli, Giada Minelli, Lorenza Nisticò, Graziano Onder, Lucia Palmisano, Daniele Petrone, Patrizio Pezzotti, Flavia Pricci, Ornella Punzo, Vincenzo Puro, Federica Quarata, Valeria Raparelli, Flavia Riccardo, Simone Rocchetto, Paolo Salerno, Giulia Sarti, Debora Serra, Matteo Spuri, Paola Stefanelli, Marco Tallon, Manuela Tamburo De Bella, Dorina Tiple, Marco Toccaceli Blasi, Federica Trentin, Brigid Unim, Luana Vaianella, Nicola Vanacore, Maria Fenicia Vescio, Emanuele Rocco Villani, Silvio Brusafferro.